



An Additive Decomposition Theorem for Analytic Functions

Author(s): Donald E. Myers

Source: *Proceedings of the American Mathematical Society*, Vol. 27, No. 3 (Mar., 1971), pp. 525-526

Published by: American Mathematical Society

Stable URL: <http://www.jstor.org/stable/2036489>

Accessed: 17/11/2008 16:54

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at <http://www.jstor.org/page/info/about/policies/terms.jsp>. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.

Please contact the publisher regarding any further use of this work. Publisher contact information may be obtained at <http://www.jstor.org/action/showPublisher?publisherCode=ams>.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

JSTOR is a not-for-profit organization founded in 1995 to build trusted digital archives for scholarship. We work with the scholarly community to preserve their work and the materials they rely upon, and to build a common research platform that promotes the discovery and use of these resources. For more information about JSTOR, please contact support@jstor.org.



American Mathematical Society is collaborating with JSTOR to digitize, preserve and extend access to *Proceedings of the American Mathematical Society*.

<http://www.jstor.org>

AN ADDITIVE DECOMPOSITION THEOREM FOR ANALYTIC FUNCTIONS

DONALD E. MYERS

ABSTRACT. Kraut, Busenberg, and Hall have shown that the uniqueness of the additive decomposition for functions of several variables given by Bochner is due to the asymptotic properties of the summands. In this paper, the uniqueness theorem is used to extend the class of functions possessing unique decompositions.

Recently Kraut, Busenberg and Hall [2] have observed that the asserted uniqueness for additive decompositions of holomorphic functions on tubes in complex n -space as given by Bochner [1] is due to the asymptotic decay of the summands. In this note we will show that [1], [2] actually provide somewhat more general results than are stated. Following [1], a tube T in C^n is of the form $\{z \mid z = (z_1, \dots, z_n), \sigma_i < \Re(z_i) < \mathfrak{I}_i\}$. A tube is said to be octant shaped if for each i either $\sigma_i = -\infty$ or $\mathfrak{I}_i = +\infty$ but not both. As observed in [2], Bochner's hypothesis for a unique decomposition is sufficient and the following theorem shows that it is not necessary.

THEOREM. *Let f be analytic in a tube T and suppose there exists a point $\alpha = (\alpha_1, \dots, \alpha_n)$ not in the closure of the base of T and an n -tuple $\beta = (\beta_1, \dots, \beta_n)$, $0 < \beta_i < 1$, such that*

$$|f(z)| \prod_{i=1}^n |z_i - \alpha_i|^{\beta_i} < M$$

for some positive constant M . Then there exist unique functions f_1, \dots, f_{2^n} each analytic in an octant shaped tube such that $f = f_1 + \dots + f_{2^n}$ in T .

PROOF. Clearly $g(z) = f(z) / \prod_{i=1}^n (z_i - \alpha_i)$ satisfies Bochner's condition of being square integrable with respect to the imaginary part for each z_i . Then $g(z)$ has a decomposition $g = g_1 + \dots + g_{2^n}$ and hence $f = f_1 + \dots + f_{2^n}$ where $f_j(z) = g_j(z) \prod_{i=1}^n (z_i - \alpha_i)$. It is only necessary then to show that each $f_j(z)$ is unique, i.e. independent of $\alpha = (\alpha_1, \dots, \alpha_n)$. By Theorem 1 [2], it is sufficient to show each f_j is

Presented to the Society, November 21, 1970 under the title *An additive decomposition theorem*; received by the editors June 24, 1970.

AMS 1970 subject classifications. Primary 32A07, 32A25; Secondary 30A84, 30A86.

Key words and phrases. Additive decomposition, tube, domain uniqueness.

Copyright © 1971, American Mathematical Society

bounded in the octant-shaped tube T_j and $f_j \rightarrow 0$ as any one of the $x_j \rightarrow \pm \infty$ in T_j .

Since the construction of the summands, as given by Bochner, is typified by the case $n=1$ we will verify the result for $n=1$ and indicate the general case. Let the base of T be $\{x \mid \sigma < x < \mathfrak{I}\}$. Then $T_1 = \{z \mid \operatorname{Re} z > \sigma\}$, $T_2 = \{z \mid \operatorname{Re} z < \mathfrak{I}\}$.

$$\begin{aligned} f(z) &= (z - \alpha)g(z) = \frac{(z - \alpha)}{2\Pi i} \int_C \frac{[f(w)/(w - \alpha)]}{w - z} dw \\ &= \frac{-1}{2\Pi i} \int_C \frac{f(w)}{w - \alpha} dw + \frac{1}{2\Pi i} \int_C \frac{f(w)}{w - z} dw \\ &= \frac{1}{2\Pi} \int_{-\infty}^{\infty} \frac{f(\mathfrak{I} + iv)}{\mathfrak{I} + iv - z} dv + \frac{1}{2\Pi} \int_{\infty}^{-\infty} \frac{f(\sigma + iv)}{\sigma + iv - z} idv \\ &= f_2(z) + f_1(z), \end{aligned}$$

since

$$\frac{1}{2\Pi i} \int_{\mathfrak{I}}^{\sigma} \frac{f(u + it)}{u + it - z} du \rightarrow 0 \quad \text{and} \quad \frac{1}{2\Pi i} \int_{\sigma}^{\mathfrak{I}} \frac{f(u - it)}{u - it - z} du \rightarrow 0$$

as $t \rightarrow \infty$ and $\sigma < \operatorname{Re} z < \mathfrak{I}$.

$$\begin{aligned} |f_2(z)| &= \frac{1}{2\Pi} \left| \int_{-\infty}^{\infty} \frac{f(\mathfrak{I} + iv)}{\mathfrak{I} + iv - z} dv \right| \\ &\leq \frac{1}{2\Pi} \int_{-\infty}^{\infty} \frac{M}{|\mathfrak{I} + iv - z| |\mathfrak{I} + iv - \alpha|^{\beta}} dv \\ &\leq \frac{1}{2\Pi} \int_{-\infty}^{\infty} \frac{M}{|v|^{1+\beta}} dv < \infty \end{aligned}$$

for $\operatorname{Re} z < \mathfrak{I}$, so that f_2 is bounded and similarly for f_1 . Furthermore, as $x \rightarrow \pm \infty$, f_1 and f_2 respectively tend to zero. The general case merely requires iterations.

REFERENCES

1. S. Bochner, *Bounded analytic functions in several variables and multiple Laplace integrals*, Amer. J. Math. **59** (1937), 732.
2. E. Kraut, S. Busenberg and W. Hall, *On an additive decomposition of functions several complex variables*, Bull. Amer. Math. Soc. **74** (1968), 372-374. MR **36** #2826.

UNIVERSITY OF ARIZONA, TUCSON, ARIZONA 85721