

Lecture Note 6: Special Distributions continued (CB 3.3-3.4)

1. Gamma Distribution The Gamma distribution with parameters $\alpha > 0$ and $\beta > 0$ is

$$f_X(x) = \frac{x^{\alpha-1} e^{-\frac{x}{\beta}}}{\Gamma(\alpha)\beta^\alpha},$$

for $x > 0$ and 0 elsewhere. Recall that the gamma function is defined as

$$\Gamma(\alpha) = \int_0^\infty t^{\alpha-1} e^{-t} dt.$$

It has the properties that

$$\Gamma(\alpha + 1) = \alpha \cdot \Gamma(\alpha), \quad \Gamma(1) = \int_0^\infty e^{-t} dt = 1, \quad \Gamma(1/2) = \sqrt{\pi},$$

with the first two implying that

$$\Gamma(n) = (n - 1)!$$

for integer n . Setting $\beta = 1/\lambda$ and $\alpha = 1$ we get the exponential distribution with arrival rate λ , so the exponential distribution is a special case of the Gamma.

The moment-generating function of the Gamma distribution is

$$M_X(t) = \frac{1}{(1 - \beta t)^\alpha},$$

and the cumulant generating function is

$$K_X(t) = -\alpha \ln(1 - \beta t).$$

The mean and variance are $\alpha\beta$ and $\alpha\beta^2$ respectively.

The Gamma distribution can be motivated in a similar way to the waiting-time motivation of the exponential distribution. In the previous lecture note we showed that the exponential distribution can be thought of as the waiting time until the first occurrence of an event. In the same way, the Gamma distribution represents the waiting time until the r th event. Let X be this waiting time. Then

$$F_X(x) = Pr(X \leq x) = 1 - Pr(\text{fewer than } r \text{ events in interval } [0, x]).$$

The last probability is equal to the probability that a Poisson random variable with arrival rate λx is less than r :

$$F_X(x) = 1 - \sum_{k=0}^{r-1} \frac{e^{-\lambda x} (\lambda x)^k}{k!}.$$

(Note that if we set $r = 1$, we have $F_X(x) = 1 - \exp(-\lambda x)$, the cumulative distribution function for the exponential distribution, the waiting time until the first event.) Take the derivative with respect to x to get the probability density function:

$$f_X(x) = \sum_{k=0}^{r-1} \frac{\lambda^{k+1} x^k e^{-\lambda x}}{k!} - \sum_{k=1}^{r-1} \frac{\lambda^k x^{k-1} e^{-\lambda x}}{(k-1)!}.$$

Note that the second summand is only from $k = 1$ to $r - 1$, not from $k = 0$ to $r - 1$. Changing the second summation from $k = 1$ to $k = r - 1$ to the summation from $k = 0$ to $k = r - 2$, we can write this as

$$\begin{aligned} f_X(x) &= \sum_{k=0}^{r-1} \frac{\lambda^{k+1} x^k e^{-\lambda x}}{k!} - \sum_{k=0}^{r-2} \frac{\lambda^{k+1} x^k e^{-\lambda x}}{k!} \\ &= \frac{\lambda^r}{(r-1)!} x^{r-1} e^{-\lambda x}. \end{aligned}$$

This is a Gamma distribution with parameters $\alpha = r$ and $\beta = 1/\lambda$. When α is not an integer, we cannot use the interpretation as the waiting time until the α th event, but the probability density is still well defined, and the extra flexibility is often useful for modelling purposes.

Since the Gamma distribution can be interpreted as the waiting time for the r th event, the following result makes sense. Suppose that Y_1, \dots, Y_r are independent exponential random variables with parameter λ . Then $X = \sum_{i=1}^r Y_i$ is a Gamma random variable with parameters $\alpha = r$ and $\beta = 1/\lambda$. More generally, if Y_1, \dots, Y_r are independent Gamma random variables with parameters α_i and β , then $X = \sum Y_i$ is Gamma with parameters $\alpha = \sum_{i=1}^r \alpha_i$ and β .

2. Chi-squared Distribution A special case of the Gamma distribution is the Chi-squared distribution. Take $\alpha = k/2$, where k is a positive integer, and $\beta = 2$, we have a Chi-squared distribution with degrees of freedom equal to k . Its pdf is

$$f_X(x) = \frac{x^{k/2-1} e^{-x/2}}{\Gamma(k/2) 2^{k/2}},$$

for x positive. Its moment generating function is

$$M_X(t) = \frac{1}{(1-2t)^{k/2}},$$

and its mean and variance are k and $2k$ respectively.

3. Normal Distribution One of the most important distributions is the normal distribution. It does not have as easy a motivation as some of the other distributions, but it is of fundamental importance as an approximation to a large number of statistics through the central limit theorem. A random variable X has a normal distribution with parameters μ and σ^2 , denoted by $\mathcal{N}(\mu, \sigma^2)$, if its pdf is

$$f_X(x) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{(x-\mu)^2}{2\sigma^2}\right),$$

for $-\infty < x < \infty$, with the parameter space $-\infty < \mu < \infty$ and $\sigma^2 > 0$. First consider the moment generating function. Its derivation relies on a trick we have used before, namely using the fact that the pdf and pmf respectively integrate and add up to one:

$$M_X(t) = \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(xt - \frac{(x-\mu)^2}{2\sigma^2}\right) dx$$

$$\begin{aligned}
&= \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{(x-\mu)^2 - 2\sigma^2 xt}{2\sigma^2}\right) dx \\
&= \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(t^2\sigma^2/2 + \mu t - \frac{(x-\mu-\sigma^2 t)^2}{2\sigma^2}\right) dx \\
&= \exp(\mu t + \sigma^2 t^2/2) \cdot \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{(x-\mu-\sigma^2 t)^2}{2\sigma^2}\right) dx \\
&= \exp(\mu t + \sigma^2 t^2/2).
\end{aligned}$$

The cumulant generating function is $K_X(t) = \mu t + \sigma^2 t^2/2$ and hence the mean is μ and the variance σ^2 .

One of the most important properties of the normal distribution is that linear transformations of normal random variables are also normally distributed. Consider a random variable X with a $\mathcal{N}(\mu, \sigma^2)$ distribution, and consider the transformation $Y = a + bX$. Then, through the moment generating function,

$$\begin{aligned}
M_Y(t) &= E(e^{tY}) = E(e^{(a+bX)t}) = e^{at} \cdot E(e^{(bt)X}) \\
&= e^{ta} \cdot M_X(bt) = \exp(at + \mu bt + \sigma^2 b^2 t^2/2) = \exp((a + b\mu)t + b^2 \sigma^2 t^2/2) \\
&= \exp(\tilde{\mu}t + \tilde{\sigma}^2 t^2/2).
\end{aligned}$$

Hence Y has a normal distribution with mean $\tilde{\mu} = a + b\mu$ and variance $\tilde{\sigma}^2 = b^2 \sigma^2$. A particularly useful transformation is that from X to $Y = (X - \mu)/\sigma$. Then $Y \sim \mathcal{N}(0, 1)$, the standard normal distribution with mean zero and unit variance.

Finally, there is a close connection between the normal distribution and the Chi-squared distribution. If X has a standard normal distribution $\mathcal{N}(0, 1)$, then $Y = X^2$ has a Chi-squared distribution with degrees of freedom equal to one. One argument goes as follows: The moment generating function of Y is

$$\begin{aligned}
M_Y(t) &= E(e^{tY}) = E(e^{tX^2}) = \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}} \exp\left(-\frac{1}{2}x^2 + tx^2\right) dx \\
&= \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}} \exp\left(-\frac{1}{2/(1+2t)}x^2\right) dx \\
&= \sqrt{1/(1-2t)} \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi/(1-2t)}} \exp\left(-\frac{1}{2/(1-2t)}x^2\right) dx = \frac{1}{(1-2t)^{1/2}}.
\end{aligned}$$

This is the mgf for a Chi-squared distribution with degrees of freedom equal to one.

4. Cauchy Distribution A random variable X has a Cauchy distribution centered around θ if it has pdf

$$f_X(x) = \frac{1}{\pi} \cdot \frac{1}{1 + (x - \theta)^2},$$

for $-\infty < x < \infty$. This distribution has no moments. The median and mode are both equal to θ . The pdf looks very similar to the pdf for the normal distribution but it has thicker tails. It is often used for modelling variables with high kurtosis, that is, which infrequently take on extremely large values, such as stock prices. An interesting property of the Cauchy distribution is that if the sequence of independent random variables X_1, X_2, \dots, X_n all have the same Cauchy distribution centered around θ , then the average $\bar{X} = \sum_{i=1}^n X_i/n$ also has that same Cauchy distribution centered around θ . In other words, laws of large numbers will be seen not to apply to distributions like Cauchy distributions.

5. Beta Distribution. Suppose two independent random variables Y_1 and Y_2 have Gamma distributions with parameters α and 1 and β and 1. Then the ratio $X = Y_1/(Y_1 + Y_2)$ has a Beta distribution with parameters α and β . The pdf is

$$f_X(x) = \frac{\Gamma(\alpha + \beta)}{\Gamma(\alpha) \cdot \Gamma(\beta)} x^{\alpha-1} \cdot (1-x)^{\beta-1},$$

for $0 < x < 1$ and zero elsewhere. The mean and variance are $\alpha/(\alpha + \beta)$ and $\alpha\beta/((\alpha + \beta)^2(\alpha + \beta + 1))$ respectively. This is a very useful distribution for modelling variables that take on values in the unit interval.

6. t-distribution. Let X have a standard normal distribution (that is, a normal distribution with mean zero and unit variance). Let W have a Chi-squared distribution with r degrees of freedom, and let X and W be independent. Then $Y = X/\sqrt{(W/r)}$ has a t-distribution with degrees of freedom equal to r . The probability density function of Y is:

$$f_Y(y) = \frac{\Gamma((r+1)/2)}{\sqrt{\pi r} \Gamma(r/2)} \cdot \frac{1}{(1+y^2/r)^{(r+1)/2}},$$

for $-\infty < y < \infty$. The t-distribution has thicker tails than the normal distribution. For $r = 1$, it is the Cauchy distribution with $\theta = 0$.

As the degrees of freedom gets larger, then the t-distribution gets closer to the normal distribution. (Note that $E[W/r] = 1$, and $V(W/r) = 2/r$, so that as $r \rightarrow \infty$, the mean of the denominator stays at one, but the variance goes to zero.)

7. F-distribution. Suppose that V and W are independent Chi-squared random variables with degrees of freedom equal to r_1 and r_2 respectively. Then $Y = (V/r_1)/(W/r_2)$ has an F-distribution with degrees of freedom equal to r_1 and r_2 . The probability density function is

$$f_Y(y) = \frac{\Gamma((r_1 + r_2)/2)(r_1/r_2)^{r_1/2}}{\Gamma(r_1/2)\Gamma(r_2/2)} \cdot \frac{y^{r_1/2-1}}{(1+wr_1/r_2)^{(r_1+r_2)/2}},$$

for positive values of y , and zero elsewhere.

Exponential Families

Finally we look at a class of families, the exponential family of distributions.

Definition 1 A family of pdf's (or pmf's) is an exponential family if it can be written as

$$f_X(x|\theta) = h(x) \cdot c(\theta) \exp\left(\sum_{i=1}^k w_i(\theta) \cdot t_i(x)\right),$$

for $-\infty < x < \infty$.

The key feature of this representation is that we can factor the probability density or mass function into functions of the parameters θ and functions of the variable x . If we parametrize this as

$$f_X(x|\eta) = h(x) \cdot c(\eta) \cdot \exp\left(\sum_{i=1}^k \eta_i \cdot t_i(x)\right),$$

for $-\infty < x < \infty$, we refer to η as the natural parameters. Many special results follow for distributions that belong to the exponential family, as we shall see later. First we show that this family includes most of the distributions we have looked at so far:

1. Binomial distribution:

$$f_X(x) = \binom{n}{x} \cdot p^x \cdot (1-p)^{(n-x)},$$

for $x = 0, 1, 2, \dots, n$, and zero otherwise. This can be written as

$$f_X(x) = 1\{x \in \{0, 1, 2, \dots, n\}\} \cdot \binom{n}{x} \cdot (1-p)^n \cdot \exp(x \ln p / (1-p)),$$

so, $k = 1$, and

$$h(x) = 1\{x \in \{0, 1, 2, \dots, n\}\} \cdot \binom{n}{x},$$

$$c(p) = (1-p)^n,$$

$$w_1(p) = \ln(p/(1-p)),$$

$$t_1(x) = x.$$

(Recall that $1\{\cdot\}$ is the indicator function, equal to one if the condition inside the brackets is true and equal to zero otherwise.)

2. Poisson distribution:

$$f_X(x) = 1\{x \in \{0, 1, 2, \dots\}\} \cdot (1/x!) \cdot \exp(-\lambda) \exp(x \ln \lambda).$$

3. Exponential distribution:

$$f_X(x) = 1\{x > 0\} \cdot \lambda \cdot \exp(-x\lambda).$$

4. Normal distribution:

$$f_X(x) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp(\mu^2/(2\sigma^2)) \cdot \exp(-x^2/(2\sigma^2) + x\mu/\sigma^2).$$

A distribution that does not fit in the exponential family is the uniform distribution:

$$f_X(x) = 1\{a < x < b\} \cdot \frac{1}{b-a}.$$

We cannot factor the indicator function $1\{a < x < b\}$ into a function of the parameters and a function of the variable.